



Lec 2a: Neural Functioning



The Cell




Membrane molecules

Protein molecules

Double Lipid Membrane



**Various Functions -
(More to come)**



Neurons & Glia



Some of the many functions of **GLIA CELLS**




(a) Central nervous system


1/10 size
of a neuron
10X as many
50% of brain
by weight

< **Myelination**

The Neuron



The Principal Structures or Regions of a Multipolar Neuron



The IONS

Charged particles
w/extra electrons (-) or fewer electrons (+)



Sodium



Potassium



Chloride



Calcium

The Nerve Impulse

To understand how Neurons “communicate”
we first need to recognize that

Nature seeks a Balance . . .



Any GRADIENT (inequality)
between chemicals inside vs. outside cell will,
if allowed, tend toward an equilibrium...

Concentration Gradient


Electrical Gradient

Concentration Gradient




Membrane molecules

Protein molecules



Ion Gates

Neuron membrane



Inside Cell

Double Lipid Membrane

Ion Gates

Outside Cell


CONCENTRATION Gradient

Higher Concentration




Lower Concentration

via Diffusion




CONCENTRATION Gradient



via Diffusion

CONCENTRATION Gradient


Lower Concentration



Higher Concentration

via Diffusion


CONCENTRATION Gradient



Equilibrium


via Diffusion

Electrical Gradient




Identical charges
REPEL

ELECTRICAL Gradient




via Electrostatic Pressure

ELECTRICAL Gradient




via Electrostatic Pressure


Equilibrium = NO Potential




Equilibrium = NO Potential




Let's look at ion conditions in an actual neuron...



Instead of being at Equilibrium, a "Resting" Neuron is
HIGHLY POLARIZED




It reaches this **HIGHLY POLARIZED** state by moving
3 Na+ Out for every 2 K+ In




Accomplished by Energy-Requiring **Na+/K+ PUMP**

Then all gates are locked = "Resting Potential"




Giant Squid Axons



Unlike most neurons,
those of the Giant Squid are
actually visible




Compare difference
between charge
INSIDE vs. OUTSIDE
the axon




Resting Potential MNEMONIC

The Resting Cell
is Polarized,
more positive QUT
than it is INSIDE

Resting Potential
-70 mV



The AXON




Axon
Hillock

Ion
Gates

Terminal
Button


Vesicles of
Neurotransmitter

The Resting Potential




Cell is ready to FIRE!

The Action Potential




A change in the polarity of the neuron
(we'll explain how, later) initiates an
ACTION POTENTIAL

The Action Potential




Na⁺ Gates at Hillock Open, Na⁺ Enters Cell

The Action Potential




Na^+ Gates at Hillock Open, Na^+ Enters Cell

The Action Potential



This local polarity change causes the next Na^+ Gates to open & Na^+ Enters Cell


The Action Potential



This local polarity change causes the next Na+ Gates to open & Na+ Enters Cell . . .


Then the previous Na+ Gates Close

The Action Potential




Then K⁺ Gates at Hillock Open, K⁺ Exits Cell

The Action Potential




Then K⁺ Gates at Hillock Open, K⁺ Exits Cell

The Action Potential




When K⁺ exits, creates a local re-polarization to -50mV
(once again, less positive inside)

The Action Potential




Next Na⁺ Gates Open, Na⁺ Enters Cell. . .

The Action Potential




Next Na⁺ Gates Open, Na⁺ Enters Cell. . .
then previous Na⁺ Gates Close

The Action Potential




Then the next K⁺ Gates Open, K⁺ Exits Cell. . .

The Action Potential




Then the next K+ Gates Open, K+ Exits Cell. . .
then previous K+ Gates Close

The Action Potential




Next Na+ Gates Open, Na+ Enters Cell . . .

The Action Potential




Next Na⁺ Gates Open, Na⁺ Enters Cell . . .
then previous Na⁺ Gates Close

The Action Potential




Next Na+ Gates Open, Na+ Enters Cell . . .
then previous Na+ Gates Close

The Action Potential




Next K⁺ Gates Open, K⁺ Exits Cell . . .

The Action Potential



Next K⁺ Gates Open, K⁺ Exits Cell . . .


The Action Potential



Next K⁺ Gates Open, K⁺ Exits Cell . . .


then previous K⁺ Gates Close

The Action Potential




When "Spike" of Depolarization reaches Terminal. . .

The Action Potential




When "Spike" of Depolarization reaches Terminal,
CA++ enters cell & Neurotransmitter released –

The Action Potential




When "Spike" of Depolarization reaches Terminal,
CA⁺⁺ enters cell & Neurotransmitter released –

The Cell Fires!




“Spike” of an
Action Potential
+50 mV

Restoring the Resting Potential



Final K⁺ gates open, and K⁺ exits cell


Restoring the Resting Potential




Polarity is back to negative, as it was initially, BUT - -


K⁺ and Na⁺ are wrong side of membrane!

Restoring the Resting Potential



Energy requiring 
Sodium–Potassium Pump

Restoring the Resting Potential




Energy requiring



Sodium–Potassium Pump

Takes in 2K⁺ for every 3⁺ Na⁺ it puts out

Restoring the Resting Potential




Energy requiring




Sodium–Potassium Pump

Takes in 2K^+ for every 3Na^+ it puts out

Restoring the Resting Potential



Energy requiring




Sodium–Potassium Pump

Takes in 2K⁺ for every 3Na⁺ it puts out

Calcium Pump

Ejects Ca⁺⁺ from Terminal


Restoring the Resting Potential



While Resting Potential is being restored,
cell can NOT fire


Refractory Period

Restoring the Resting Potential



Once Resting Potential is fully restored cell is

ready to FIRE!



MNEMONICS for Resting Potential

When Ions of Sodium want to come in,
what does the Resting Cell say?

Na+, Na+, Na+

This is because what minority is locked inside?


K+ K+ K+

(Actually **K+K+** for each **Na+Na+Na+**)


Action Potential = "All or Nothing"

Action Potential results in same release of NT
regardless of intensity of input

(as long as "Threshold for Firing" is crossed)





Every spike is the same



But message can vary
Temporally!

Myelinization






Glia Cells,
wrapped around the axon,
w/gaps between called
"Nodes of Ranvier"

Electrical Conduction


Ionic Conduction
(Ions flow across membrane)

(Electricity flows thru axon
under "insulation")



Slow, but stays strong

VERY fast,
but decays over distance




Myelinization

Glia Cells,
wrapped around the axon,
w/gaps between called
"Nodes of Ranvier"

Electrical Conduction

(Electricity flows thru axon
under "insulation")

Ionic Conduction
(Ions flow across membrane)




So, overall, myelinated axons show
"**Saltatory**" – or "jumping" - Conduction

Multiple Sclerosis




Multiple Sclerosis

How Multiple Sclerosis Works



Multiple Sclerosis



No ion gates under the myelin so signal does not propagate

Graded Potentials

Not all Neurons show "Action Potentials"

Cells that fire "Graded Potentials" may release
MORE or **LESS** neurotransmitter

e.g. "Hair cells"
Auditory receptors

Soft sound,
cilia move a
little



A little
neurotransmitter
is released

NT
NT

Loud sound,
cilia move
a lot



A lot of
neurotransmitter
is released

NT NT NT
NT NT NT
NT NT NT